

GLUEBALL-SIGMA IDEA AND PRC-USA
(IHEP-CMU) COLABORATION

SCALAR GLUEBALLS AND SIGMA DECAY

THE SIGMA-GLUEBALL MODEL

THE POMERON AND THE SIGMA-GLUEBALL

PROTON-PROTON PRODUCTION OF SIGMAS

UPSILON DECAYS TO SIGMAS

HYBRID QUARKONIUM AND THE SIGMA-GLUEBALL

D-DECAYS TO SIGMAS

NPQCD AND THE $\rho - \pi$ PUZZLE

LEONARD KISSLINGER

IHEP PRC-USA COLLABORATION MEETING 2006

IHEP-CMU THEORY TEAM

IHEP MEMBERS

WEI-XING MA

PENGNIAN SHEN

FENG-KUN GUO

LIJUAN ZHOU

CMU MEMBERS

LEONARD KISSLINGER


SEAMUS RIORDAN

DIANA SEYMOUR

SAMEER WALAWALKAR

SCALARS: MESON/GLUEBALL CANDIDATES

THE FOLLOWING SCALARS WERE DISCUSSED AT
CHARM 2006 AS MESON/GLUEBALL CANDIDATES



THE SIGMA [$I=0$, $L=0$, $\pi - \pi$ RESONANCE]

Zou-Bugg, Phys. Rev. D50, 591 (1994)




FIG. 1. The isoscalar S -wave phase shift δ_0 for $\pi\pi$ scattering [18–20]. The dashed line includes only the contribution from the t -channel ρ exchange. The dot-dashed line includes the contribution of the s -channel resonance $f_0(975)$ in addition. The solid line further includes the contribution from the t -channel $f_2(1275)$ exchange.

The $\pi^+\pi^-\pi^+\pi^-$ spectrum of $J/\psi \rightarrow \gamma(\pi^0)4\pi$




Figure 11

June 1997 L.Y. DONG / BES CONF97

↑
1.5 ↑
1.75 ↑
2.1

Decay Modes Used in the Final Fit

L is the orbital angular momentum between the two final state particles following the decay of the resonance

J^P	Mass	channel	L or $^{2S+1}L_2$
2^+	$f_2(1270)$	$\rho\rho$	1D_2 3D_2
0^-	$\eta(1420)$	$\rho\rho$	$L = 1$
0^+	$f_0(1500)$	$\sigma\sigma$	$L = 0$
2^+	$f_2(1560)$	$\rho\rho$	5S_2 1D_2 3D_2
2^+	$f_2(1710)$	$\rho\rho$	5S_2
0^+	$f_0(1750)$	$\sigma\sigma$	$L = 0$
2^+	$f_2(2010)$	$\sigma\sigma$	1D_2 3D_2
		$\rho\rho$	1D_2 3D_2
		$f_2(1270)\sigma$	5S_2
0^+	$f_0(2100)$	$\sigma\sigma$	$L = 0$
2^+	$f_2(2220)$	$\sigma\sigma$	5S_2 3D_2
		$\rho\rho$	5S_2 3D_2
		$f_2(1270)\sigma$	5S_2

Table 2

June 1997 L.Y. DONG / BES CONF97

SCALAR GLUEBALLS AND SIGMA DECAY

The $f_o(1500)$ and $f_o(1710)$ are glueball candidates

The 4-pion channel dominates $f_o(1500)$ and $f_o(1710)$ decay.

From the BES analysis:

$$\frac{Br(f_o(1500) \rightarrow 4\pi)}{Br(f_o(1500) \rightarrow 2\pi)} = 3.3 \pm 0.8$$

The 4-pion decay from the $f_o(1500)$ decay, from the BES analysis, is dominated by $2\text{-}\sigma$ decay, as is the $f_o(1710)$ (L.Y. Dong/BES CONF97)

This observation was an important motivation for the scalar Glueball-Sigma picture

SCALAR MESONS/GLUEBALLS VIA QCD SUM RULES

L.Kisslinger, J.Gardner, C.Vanderstraeten, PLB410, 1 (97)


L.Kisslinger, M.Johnson, PLB523, 127 (2001)

Scalar glueball, scalar meson, and mixed glueball meson operators

$$J^G(x) = \alpha_s G^2$$

$$J^m(x) = \frac{1}{2}(\bar{u}(x)u(x) - \bar{d}(x)d(x))$$

$$J_{0^{++}} = \beta M_o J_m + (1 - |\beta|) J_G$$



Glueball-meson coupling (theorem: Novikov et al, NPB191)

$$\int dx T[J^G(x)J^m(0)] \simeq -\frac{32}{9} \langle \bar{q}q \rangle$$

RESULTS


80% scalar glueball at 1500 MeV- $f_o(1500)$

80% scalar meson at 1350 MeV- $f_o(1370)$

Light Scalar Glueball 400-600 MeV-The Glueball-Sigma

SIGMA PRODUCTION IN PROTON-PROTON COLLISIONS

L.Kisslinger, W-x Ma, and P. Shen, PRD71, 094021 (2005)




$A^{pp\sigma}$ = amplitude for σ production in p-p collisions

$$A^{pp\sigma} \simeq V(t_1)\bar{D}_\sigma^P(t_1, t_2, s)V(t_2),$$

where $V(t)$ =pomeron-nucleon vertex and \bar{D}_σ^P is the Pomeron propagator with a σ emission. [See figure]

Ratio of $pp \rightarrow pp\sigma$ to elastic $pp \rightarrow pp$ scattering:



QUARKONIUM STATES: MANY NEW $Q\bar{Q}$ STATES FOUND
BY CLEO, BES, BaBar, Belle (FPCP 2006, Vancouver, BC)

Recent Effective Hadronic Hamiltonian calculations of quarkonium states by PRC theorists:

F-K. Guo, P-N. Shen, H-C. Chiang, and R-G. Ping

Nucl. Phys. A761, 269 (2005)

hep-ph/0601120

hep-ph/0603072

hep-ph/0603117

F-K. Guo, P-N. Shen, H-C. Chiang, R-G. Ping, B-S. Zou

hep-ph/0509050

IHEP-CMU theoretical studies being planned on $Q\bar{Q}$ states:

Further use of hadronic models to predict quarkonium states

P-N. Shen, F-K. Guo and collaborators

Bethe-Salpeter and Dyson equations will be used to study amplitudes:

W-x. Ma and Zhou Lijian and collaborators





QCD sum rules will be used for NPQCD investigations of new states:

All theorists will collaborate.

CHARMONIUM TRANSITIONS WITH SIGMA EMISSION

EXAMPLE: THE σ POLE IN $J/\Psi \rightarrow \omega\pi^+\pi^-$

M. Abilkim et. al (BES Collaboration), Phys. Lett. B598, 149 (2004)




UPSILON TRANSITIONS WITH SIGMA EMISSION




$\Delta n=2$ CONJECTURE: H.Vogel at FPCP 2006, Vancouver, BC

Belle: hep-ex/0512034
 BaBar: talk at QCD Moriond'06
 CLEO: very preliminary




Belle 2S-1S




Belle 4S-1S



UPSILON TRANSITIONS WITH SIGMA EMISSION (continued)



POSSIBLE SIGMA-GLUBALL TREATMENTS OF σ PRODUCTION IN QUARKONIUM TRANSITIONS




Studies by L. Kisslinger, Seamus Reardon and Diana Seymour, and collaborators are being planned.

HYBRID CHARMONIUM AND SIGMA-GLUEBALL


Example of possible hybrid baryon: the Roper

L. Kisslinger and Z. Li, Phys. Rev. D51, R5986 (1995)
Phys. Lett. B445, 271 (1999)



HYBRID QUARKONIUM STATES (see, e.g., S Godfrey, FPCP 2006, Vancouver, BC)

Expect sigma decay:



Our IHEP-CMU group expects to investigate such hybrid quarkonium states via sigma decays.

NEW CHARMONIUM HYBRID CANDIDATES

As discussed by G. Rong (IHEP) at CHARM 2006:

$$\psi(3770) \rightarrow J/\psi \pi^+ \pi^- (\sigma)$$

and as widely discussed by a number of speakers at CHARM 2006 the

$$Y(4260)$$

does not fit the $c\bar{c}$ spectrum of successful models.

The $Y(4260)$ is a hot new state, and experimental investigations of its properties are planned.

Our IHEP-CMU theory group will study these as $c\bar{c}g$ hybrids and predict decay rates.

D-DECAYS TO SIGMAS

“Experimental Evidence for a Light and Broad Scalar resonance in $D^+ \rightarrow \pi^- \pi^+ \pi^+$ Decay”, E.M. Aitala et al (Fermilab E791 Collaboration), Phys. Rev. Lett. 86, 770 (2001)




FIG. 2. s_{12} and s_{13} projections for data (error bars) and fast MC (solid line). The shaded area is the background distribution, (a) solution with the Fit-1, and (b) solution with Fit 2.

D-DECAYS TO SIGMAS

“Experimental Evidence for a Light and Broad Scalar resonance in $D^+ \rightarrow \pi^- \pi^+ \pi^+$ Decay”, E.M. Aitala et al (Fermilab E791 Collaboration), Phys. Rev. Lett. 86, 770 (2001)




FIG. 2. s_{12} and s_{13} projections for data (error bars) and fast MC (solid line). The shaded area is the background distribution, (a) solution with the Fit 1, and (b) solution with Fit 2.

PROCESSES FOR D DECAYS INTO SIGMAS

Using an effective Hamiltonian (L.Kisslinger, hep-ph/0103326) the unique skeletal diagram, with an internal W^+ is (Fig 1):




Fig. 1 D^+ decay to π^+ and ϕ or σ (a) via internal W^+ emission or (b) penguin diagram

The matrix element for decay into π^+ and σ, ϕ (called X) is

$$\langle X\pi^+ | H^{eff} | D^+ \rangle = \frac{G_F}{\sqrt{2}} \cos\theta_c \sin\theta_c C_3 \langle X | (\bar{s}s)_L | 0 \rangle \langle \pi^+ | (\bar{u}c)_L | D^+ \rangle$$

Taking the ratio, so constants drop out, and using known values of the gluon and quark condensates it was estimated that

$$\frac{\Gamma(D^+ \rightarrow \sigma\pi^+)}{\Gamma(D^+ \rightarrow \phi\pi^+)} \simeq 0.15.$$

This value is in agreement with E791 experimental results.

PROCESSES FOR D DECAYS INTO SIGMAS (continued)

Other known processes, the annihilation-hairpin and final state interactions are illustrated in Fig.2




Fig. 2 (a) Annihilation-hairpin, (b) Final state interaction

These processes are harder to evaluate, and are a subject for IHEP-CMU theoretical research

PROCESSES FOR D_s and τ DECAYS INTO SIGMAS

For D_s decays into π^+ a σ or a ϕ the spectator and annihilation diagrams are illustrated in Fig. 3




Fig. 3 (a) Spectator, (b) Annihilation

A rough evaluation (LSK hep-ph/0103326) finds the ratio of σ to ϕ rates for D_s decay are reduced in comparison with D decay, in general agreement with E791.

The skeletal diagram for $\tau^- \rightarrow \sigma$ is illustrated in Fig. 4.





Fig. 4 Diagram for $\tau^- \rightarrow \nu_\tau \pi^- \sigma$ (or scalar meson)

The IHEP-CMU theorists plan to investigate D and τ decay processes, particularly looking for σ decays.

THE $\rho - \pi$ PUZZLE FOR $c\bar{c}$ STATES

Using the lowest order perturbative diagram for $Q\bar{Q}$ decays into hadrons, illustrated below,

LOWEST ORDER DIAGRAMS FOR $Q\bar{Q}$ DECAYS



by taking ratios the wave functions at the origin cancel, and for many years it has been known that for $c\bar{c}$ decays into hadrons (h) the ratios of branching rates

$$R = \frac{B(\Psi'(c\bar{c}) \rightarrow h)}{B(J/\Psi(c\bar{c}) \rightarrow h)} = \frac{B(\Psi'(c\bar{c}) \rightarrow e^+e^-)}{B(J/\Psi(c\bar{c}) \rightarrow e^+e^-)} \simeq 0.15$$

for all hadronic decays except the $\rho - \pi$ decay. The $\rho - \pi$ decay ratio is more than an order of magnitude smaller. This is the famous $\rho - \pi$ puzzle. Many, many theoretical attempts to explain this puzzle have still left the puzzle.

The IHEP-CMU theorists will explore the sigma-gluon diagrams for nonperturbative QCD as a possible solution to this puzzle.