The Analysis of Decay Channels $J/\psi \to \gamma\gamma V(\rho, \phi)$

Xu Guofa

June 15, 1999

Abstract

Based on 7.8×10^6 J/ψ events collected by the BES experiment, J/ψ radiative decay channels $J/\psi \to \gamma\gamma V(\rho, \phi)$ have been studied. The $\eta(1430)$ resonance has been observed in these two channels respectively. Its mass and width are 1431 ± 17 MeV and 88 ± 28 MeV from the $J/\psi \to \gamma\gamma\rho$ mode, 1427 ± 13 MeV and 77 ± 46 MeV from the $J/\psi \to \gamma\gamma\phi$ mode.

1 Introduction

Up to now, the structure of $\iota(1440)$ is not very clear. The existence of two overlapping pseudo-scalar states has been suggested, one of which around 1410 MeV decays into both $K\overline{K}\pi$ and $\eta\pi\pi$, and the other one around 1470 MeV decays only to $K\overline{K}\pi$. It is therefore conceivable that the higher mass state is the $s\bar{s}$ member of the 2^1S_0 nonet[1]. Is the lower mass state a meson, glueball, or hybrid? It should be further investigated in different decay channels.

The radiative decay of $\iota(1440) \to \gamma\rho$ has been analysed in the process $J/\psi \to \gamma\gamma\rho(\rho \to \pi^+\pi^-)$ by Crystal-Ball[2], MarkIII[3] and DM2[4] respectively(see Tab. 1). From their analysis of the $\gamma\rho$ decay angular distribution, $\iota(1440)$ is mainly associated with a spin 0^- interpretation. However, spin 1^+ still cannot be excluded.

Table 1: Experimental results for $X(1440) \to \gamma\rho$

Mass (MeV)	Width (MeV)	Br$(J/\psi \to \gamma X)Br(X \to \gamma\rho)$ $(\times 10^{-4})$	Exp.
1390 ± 25	185^{+110}_{-80}	$1.9 \pm 0.5 \pm 0.4$	Cry. Ball
1432 ± 8	90 ± 26	$0.64 \pm 0.12 \pm 0.07$	MarkIII
1401 ± 18	174 ± 44	$0.9 \pm 0.2 \pm 0.14$	DM2

A pure $s\bar{s}$ meson can decay radiatively into $\gamma\phi$, but not $\gamma\rho$. A pure $(u\bar{u} + d\bar{d})$ meson can decay radiatively into $\gamma\rho$, but not $\gamma\phi$. A glueball can mix with both $s\bar{s}$ and $(u\bar{u} + d\bar{d})$, and therefore can decay into both $\gamma\phi$ and $\gamma\rho$. Here we report a study of $J/\psi \to \gamma\gamma\rho$ and $J/\psi \to \gamma\gamma\phi$ channels based on 7.8×10^6 J/ψ events collected by the BES experiment.

2 Event Selection

In this paper we mainly analyse the following channels :

$$J/\psi \to \gamma X(X \to \gamma V)$$

where V includes vectors ρ and ϕ; their decay modes are as follows:

$$\rho \to \pi^+\pi^-$$
$$\phi \to K^+K^-.$$

The BES detector has been fully described elsewhere [5]. For the above processes, two oppositely charged particles should be detected by the MDC (Main Drift Chamber).

The subdetector BSC (Barrel Shower Counter) of BES covers 80% of 4π solid angle, and has an energy resolution of $\delta E/E = 22\%/\sqrt{E}$. In order to exclude electronic noise, the energy deposited in BSC by each neutral particle is required to be $E_{sc} > 70MeV$. The photon is also required to be isolated from two charged tracks ($cos(\theta_{\gamma\pi(K)}) < 0.98$); this also rejects any radiative photon from charged particles. Photons are also required to come from the interaction point. We call photons passing these cuts "good" photons. The photon with maximum energy among all of those "good" photons is selected as the J/ψ radiative photon.

2.1 The event selection of $J/\psi \to \gamma\gamma\rho(\rho \to \pi^+\pi^-)$

We define P_{miss} as the total momentum of all charged particles. A cut of $P_{miss} \leq 1.25GeV/c$ is required. This criterion rejects the background channel $J/\psi \to \rho\pi^0$ effectively.

If the weight of π identified by time of flight (TOF) is greater than that of K, we assume this track to be a π . Finally at least one π track is required per event.

In order to obtain a good $J/\psi \to \gamma\gamma\pi^+\pi^-$ event, we do a kinematic fit (4C-FIT) for all four particle final states which include two photons and two charged particles. Because of the existence of pseudo-photons and photons from multi-photon backgrounds, we take the maximum energy photon and two charged particles as the first three particles, and one of the photons other than the maximum energy photon as the fourth particle in the 4C-FIT. Every photon except the maximum energy photon is treated sequentially in this 4C-FIT procedure. We select the event with the minimal $\chi^2(\chi^2_{min})$ ($\chi^2_{min} \leq 6$) as a "good" event. We call photons finally passing our χ^2 cut "true" photons(including the maximum energy photon). In order to suppress background channels which include the following final states : $2\gamma K^+K^-$, $\pi^0\pi^+\pi^-$, $\eta\pi^+\pi^-$, $3\gamma\pi^+\pi^-$, and $\gamma\pi^+\pi^-$, the event we selected above has been subjected to a 4C-FIT to all the above final states. If the χ^2 for any of these channels satisfies $\chi^2 < \chi^2_{min}$, it will be rejected as background. Another useful cut, which can effectively suppress multi-photon backgrounds, requires that the total energy of photons not accepted as "true" photons $E_{rest} \leq 130MeV$. Fig. 1 shows the invariant mass spectrum of $\pi^+\pi^-$; the ρ signal is selected as $|M_{\pi^+\pi^-} - M_\rho| \leq 0.14GeV/c^2$. The invariant mass spectrum of $\gamma\rho$ is shown in Fig. 2.

The detection efficiency is determined from a Monte Carlo simulation. After the same selections as real data, we obtain a detection efficiency of 4.8%. The systematic errors listed in Tab. 2 are determined by adding (i) the error in our event selection (6.0%), (ii) different background shapes for the BW fit (3.0%), (iii) the uncertainty of the Monte Carlo simulation

of the detection efficiency (14%), and (iv) the uncertainty in the number of produced J/ψ (10%). The systematic error in this J/ψ decay mode is 19% in total.



Figure 1: The Invariant Mass Spectrum of $\pi^+\pi^-$



Figure 2: The Invariant Mass Spectrum of $\gamma\rho$. The upper curve is the resonance contribution, and the lower one is the shape of backgrounds.

2.2 The event selection of $J/\psi \to \gamma\gamma\phi$ $(\phi \to K^+K^-)$

If the weight of K which is identified by TOF is greater than that of π, we assume this track as K. Eventually at least one K track is required. In order to suppress multi-photon backgrounds, P_{miss} should satisfy $P_{miss} \leq 1.25GeV/c$.

In order to get good $J/\psi \to \gamma\gamma K^+K^-$ events, we do a 4C-FIT for four final states. The 4C-FIT procedure is similar to that in the $J/\psi \to \gamma\gamma\pi^+\pi^-$ process. We select the event with $\chi^2_{min} \leq 25$ as a "good" event. Furthermore, if the χ^2 of the "background" channels $2\gamma\pi^+\pi^-$, γK^+K^-, and $3\gamma K^+K^-$ satisfy $\chi^2 > \chi^2_{min}$, this event is considered as a good event; otherwise, the event (with $\chi^2 < \chi^2_{min}$) is rejected as background. Fig. 3 shows the invariant mass spectrum of K^+K^- .

Table 2: The resonance parameters of $J/\psi \to \gamma X(X \to \gamma\rho)$

Mass (MeV)	Width (MeV)	$\mathrm{Br}(J/\psi \to \gamma X)Br(X \to \gamma\rho)$ $(\times 10^{-4})$	Events	Significance
961 ± 1	11 ± 3	$10.8 \pm 0.4 \pm 2.1$	269 ± 10	30.7σ
1245 ± 11	20 ± 16	$0.5 \pm 0.2 \pm 0.1$	13 ± 12	3.0σ
1431 ± 17	88 ± 28	$1.5 \pm 0.6 \pm 0.3$	38 ± 16	4.4σ



Figure 3: The Invariant Mass Spectrum of K^+K^-

By taking $M_{\gamma\gamma} \geq 0.6GeV/c^2$, the backgrounds $J/\psi \to \phi\eta$ and $J/\psi \to \phi\pi^0$ can be rejected . Requiring $|M_{K^+K^-} - M_\phi| \leq 0.02GeV/c^2$ as the ϕ signal, the final $\gamma\phi$ spectrum shown in Fig. 4 is obtained. Fig. 5 shows the γK^+K^- invariant mass spectrum of a ϕ side-band. There is no structure around 1430 MeV.

After the Monte Carlo simulation, we obtain a detection efficiency of 6.2%.

The systematic errors listed in Tab. 3 are determined in a manner similar to those in Tab. 2. They include (i) the error in our event selection (8%), (ii) different background shapes for the BW fit (11%), (iii) the uncertainty of the Monte Carlo simulation of the detection efficiency (10%), (iv) the possible backgrounds due to π/K particle identification (10%), and (v) the uncertainty in the number of produced J/ψ (10%). Finally, the systematic error in this J/ψ decay mode is 22% total.

3 Results and Discussions

First we discuss results from $J/\psi \to \gamma\gamma\rho$. MarkIII observed three resonances [3] similar to our results. The first state, with a mass of 961 MeV in Tab. 2, can be interpreted as $\eta'(958)$. Our result is compatible with the PDG branching ratio $(Br(J/\psi \to \gamma\eta'(958), \eta'(958) \to \gamma\rho) = (13.0 \pm 1.1) \times 10^{-4}$. This result includes non-resonant $\pi^+\pi^-\gamma$). The state around 1245 MeV in Tab. 2 cannot be analysed in detail, because of low statistics. MarkIII also obtained this resonance in $J/\psi \to \gamma\gamma\rho$. They interpreted it as the signal of $f_1(1285)$, but



Figure 4: The Invariant Mass Spectrum of $\gamma\phi$. The upper curve is the resonance contribution, and the lower one is the shape of backgrounds.

Table 3: The resonances parameters of $J/\psi \to \gamma X(X \to \gamma\phi)$

Mass (MeV)	Width (MeV)	Br$(J/\psi \to \gamma X)Br(X \to \gamma\phi)$ $(\times 10^{-4})$	Events	Significance
1427 ± 13	77 ± 46	$1.7 \pm 0.7 \pm 0.4$	41 ± 16	6.4σ
1833 ± 29	80 ± 78	$0.6 \pm 0.5 \pm 0.1$	14 ± 11	2.8σ

they could not reject the possibility of $\eta(1295)$ [3]. The state X(1430) will be discussed below.

Consider next results from $J/\psi \to \gamma\gamma\phi$. The signal around 1833 MeV in Tab. 3 cannot be analyzed in detail, because of low statistics .

Finally we discuss the X(1430) state observed in both $J/\psi \to \gamma\gamma\rho$ and $J/\psi \to \gamma\gamma\phi$ channels. The measured parameters of X(1430) are listed in Tab. 4.

The WA102 collaboration [6] recently gave the result:

$$\frac{f_1(1420) \to \rho^0\gamma}{f_1(1420) \to K\overline{K}\pi} < 0.02 \ (95\% \ c.l.),$$

The PDG gives the world average of the branching ratio:

$$Br(J/\psi \to \gamma f_1(1420), f_1(1420) \to K\overline{K}\pi) = (8.3 \pm 1.5) \times 10^{-4}.$$

Combining these results we obtain:

$$Br(J/\psi \to \gamma f_1(1420), f_1(1420) \to \gamma\rho) < 1.7 \times 10^{-5}.$$

Our result for X(1430) from the $J/\psi \to \gamma\gamma\rho$ channel is ten times this upper limit. The X(1430) from our $J/\psi \to \gamma\gamma\rho$ analysis should therefore be $\eta(1430)$, not $f_1(1420)$.

The WA102 collaboration [6] also gives:

$$\frac{f_1(1420) \to \phi\gamma}{f_1(1420) \to K\overline{K}\pi} = 0.003 \pm 0.001 \pm 0.001.$$



Figure 5: The ϕ Side-band

Using the PDG value of the branching ratio mentioned above, we have

$$Br(J/\psi \to \gamma f_1(1420), f_1(1420) \to \gamma\phi) = (0.249 \pm 0.068 \pm 0.099) \times 10^{-5}.$$

This is two orders of magnitude smaller than our result for the X(1430) state. So this signal from $J/\psi \to \gamma\gamma\phi$ cannot be due to $f_1(1420)$. It must be due to $\eta(1430)$ dominantly.

Table 4: Results of $\eta(\sim 1430)$ from the analysis of $J/\psi \to \gamma\gamma V$ in BES

Decay Modes	Mass (MeV)	Width (MeV)	Br$(J/\psi \to \gamma X)Br(X \to \gamma V)$ $(\times 10^{-4})$	Significance
$X \to \gamma\rho$	1431 ± 17	88 ± 28	$1.5 \pm 0.6 \pm 0.3$	4.4σ
$X \to \gamma\phi$	1427 ± 13	77 ± 46	$1.7 \pm 0.7 \pm 0.4$	6.4σ

We conclude that the X(1430) state from both $J/\psi \to \gamma\gamma\rho$ and $J/\psi \to \gamma\gamma\phi$ channels must be dominantly $\eta(1430)$. From those features of $\eta(\sim 1430)$ shown in Tab. 4, we can say that $\eta(\sim 1430)$ has a significant decay to $\gamma\phi$. With this assumption, we now discuss the relative branching ratios $r = BR[\eta(1430) \to \gamma\phi]/BR[\eta(1430) \to \gamma\rho] = 1.1 \pm 0.7$. Here, systematic errors common to both decay modes are factored out. The ratio r is proportional to phase space for each channel and also to the mean square of quark charges. The process $\eta(1430) \to \gamma V$ involves one unit of orbital angular momentum, hence phase space $\propto p^3/(p^2 + 1/R^2)$, where p is the decay momentum and R the raidus of the centrifugal barrier. Taking $R = 0.8$ fm, phase space favours the $\rho\gamma$ channel by a factor 1.7. Quark charges also favour it by a factor 2.5. Although the error on r is sizeable, our result suggests a distinctly larger $s\bar{s}$ component than $(u\bar{u} + d\bar{d})/\sqrt{2}$. That agrees with the conclusion of a previous BES publication on $J/\psi \to \gamma(K\overline{K}\pi)$ [7], where it was found that decays to $K^*(890)K$ dominate strongly over non- strange decay modes. However, further investigation of the nature of this state will be necessary and possible from a BES sample now being taken with higher statistics.

We would like to thank the staff of BEPC accelerator and IHEP Computing Center. We are greatly indebted to Profs. D.V. Bugg, B.S. Zou, H. Yu, Y.G. Xie, and Dr. J.D. Richman for valuable discussions. This work was supported by the National Natural Science Foundation and Academia Sinica.

References

[1] PDG,Eur. Phys. J. C3(1998)396
M.G.Rath et al., Phys. Rev. D40(1989)693
A.Bertin et al., Phys. Lett. B361(1995)187

[2] C.Edwards,PhD thesis,Cal.Tech.Preprint CALT-68-1165(1985)

[3] D.Coffman et al., Phys. Rev. D41(1990)1410
J.D. Richman,PhD thesis,Caltech preprint CALT-68-1231(1985)

[4] J.E.Augustin et al., Orsay preprint LAL/85-27(1985)

[5] J.Z.Bai et al., Nucl. Instrum. Methods Phys. Res. A344(1994)319
J.Z.Bai et al., Phys. Rev. Lett. 69(1992)3021

[6] D. Barberis et al., Phys. Lett. B440(1998)225

[7] J.Z. Bai et al., Phys. Lett. B440 (1998) 217