ψ' decays to Baryon Pairs

Jim Panetta

California Institute of Technology

for the BES Collaboration[†]

(May 13, 1999)

Abstract

A sample of 3.7M ψ' decays registered in the BES detector are used to study final states containing pairs of octet and decuplet baryons. We report branching ratios for $\psi' \to p\overline{p}$, $\Lambda\overline{\Lambda}$, $\Sigma^0\overline{\Sigma}^0$, $\Xi^-\overline{\Xi}^+$, $\Delta^{--}\overline{\Delta}^{++}$, $\Sigma^+(1385)\overline{\Sigma}^-(1385)$, $\Xi^0(1530)\overline{\Xi}^0(1530)$, and $\Omega^-\overline{\Omega}^+$. These results are compared to expectations based on the $SU(3)$-flavor symmetry, factorization, and perturbative QCD.

Typeset using REVTEX

I. INTRODUCTION

In the quarkonium model, the ψ' is the first radial excitation of the ^3S $c\overline{c}$ bound state. As such, its properties are expected to be relatively straight-forward to understand, at least in terms of those of the J/ψ ground state. Somewhat surprisingly, these expectations do not always hold. In particular, there is a rather dramatic anomaly associated with the ψ'.

The major puzzle in hadronic ψ decays is the large discrepancy between the decay widths for $J/\psi(1S) \to \rho\pi$ and K^*K and the corresponding widths for $\psi(2s)$ decays. These modes are expected to proceed via $\psi \to ggg$, with widths that are proportional to the square of the $c\overline{c}$ wave function at the origin, which is well determined from dilepton decays. The predicted ratio is:

$$\frac{\mathcal{B}(\psi' \to X_{had})}{\mathcal{B}(J/\psi \to X_{had})} = \left[\frac{\alpha_s(\psi')}{\alpha_s(J/\psi)}\right]^3 \frac{\mathcal{B}(\psi' \to e^+e^-)}{\mathcal{B}(J/\psi \to e^+e^-)} = 0.116 \pm 0.022,$$

where X_{had} designates any exclusive hadronic decay channel. [1] Experimentally, the $\psi(2s) \to \rho\pi$ and K^*K are reduced by over a factor of twenty from these expectations [2]. This anomaly calls into question the underlying assumption behind the theoretical predictions, namely that the ψ' is a pure $c\overline{c}$ state.

A. $\psi' \to B_i\overline{B_i}$

In the context of flavor $SU(3)$, a pure $c\overline{c}$ state is a flavor singlet and, in the limit of $SU(3)$ flavour symmetry, the phase-space-corrected reduced branching ratios to any baryon octet pair, $|M_i|^2$, where

$$|M_i|^2 = \frac{\mathcal{B}(\psi' \to B_i\overline{B_i})}{\pi p^*/\sqrt{s}}$$

(p^* is the momentum of the baryon in the ψ' rest frame), should be the same for every octet baryon, B_i. Deviations from this rule would indicate a non-$c\overline{c}$ component of the charmonium wave function. The reduced branching fractions for $J/\psi \to B_i\overline{B_i}$ decays are shown in Fig. **??**. The $SU(3)$ relation works reasonably well, although there may be some preference for the $p\overline{p}$ mode.

This relation has not been tested for the ψ', where the only relevant mode that has been measured is $p\overline{p}$, and that with rather poor precision [3], [4].

II. THIS EXPERIMENT

We report results of measurements of the branching fractions for $\psi' \to B_i\overline{B_i}$, where $B_i \in \{p, \Lambda, \Sigma^0, \Xi^-, \Delta^{++}, \Sigma^+(1385), \Xi^0(1530), \Omega^-\}$ using a sample of 3.7×10^6 ψ' events produced via e^+e^- annihilations at the BEPC collider and observed by the BES detector.

The Beijing Electron Spectrometer, BES, is a conventional cylindrical magnetic spectrometer, coaxial with the BEPC colliding e^+e^- beams [5]. A four-layer central drift chamber (CDC) surrounding the beampipe provides trigger information. Outside the CDC, a forty-layer main drift chamber (MDC) provides tracking and energy-loss (dE/dx) information on charged tracks over 85% of the total solid angle. The momentum resolution is $\sigma_p/p = 0.017\sqrt{1+p^2}$ (p in GeV/c), and the dE/dx resolution for hadron tracks is $\approx 11\%$. An array of 48 scintillation counters surrounding the MDC provides time-of-flight (TOF) information of charged tracks with a resolution of $\approx$ 450 ps for hadrons. Outside the TOF system, a 12 radiation length, lead-gas barrel shower counter (BSC), operating in self-quenching streamer mode, measures the energies of electrons and photons over $\approx 80\%$ of the total solid angle. The energy resolution is $\sigma_E/E = 0.22/\sqrt{E}$ (E in GeV), and the spatial resolutions are $\sigma_\phi = 4.5$ mrad and $\sigma_z = 4$ cm. Surrounding the BSC is a solenoidal magnet that provides a 0.4 Tesla magnetic field in the central tracking region of the detector. Three double layers of planar counters instrument the magnet flux return (MUID) and are used to identify muons of momentum greater than 0.5 GeV/c. Endcap time-of-flight and shower counters extend coverage to the forward and backward regions.

A. The ψ' data samples

The data were accumulated in two runs: an ≈ 2.2 pb^{-1} data sample accumulated between December 1993 and January 1994, and an ≈ 4.5 pb^{-1} data sample accumulated between

December 1994 and March 1995.

III. BARYON OCTET

A. $\psi' \to p\overline{p}$

The experimental signature for the decay $\psi' \to p\overline{p}$ is two back-to-back, oppositely charged tracks each with a momentum of 1.586 GeV/c. The proton typically deposits one-half or less of its 0.91 GeV kinetic energy in the BSC; the antiproton undergoes an annihilation process in the BSC approximately half the time, producing a large shower.

Major potential backgrounds are: $\psi' \to K^+K^-$, $\pi^+\pi^-$, $\mu^+\mu^-$, and e^+e^-. Each of these modes has a momentum at least 190 MeV/c greater than that of the $p\overline{p}$ channel.

1. $p\overline{p}$ selection

We select events with two and only two well reconstructed, oppositely charged tracks with good time of flight information, and which are not identified as muons by the muon system. Also $|\cos\theta|$ must be less than 0.6 for both tracks to ensure that they occur within the fiducial volume covered by the muon system. Candidate $p\overline{p}$ pairs are required to be within 1.8 degrees of collinear.

The shower counter energy deposition as a function of momentum for positively charged tracks is shown in Figure 1. The faint cluster near $p = 1.6$ GeV/c, $E_{SC} = 0.3$ GeV is the proton signal. The other features on the graph are due to Bhabhas, (large concentration at $p \approx 1.8$ GeV/c, $E_{SC} \approx 1.5$ GeV), muons (vertical stripe at $p \approx 1.8$ GeV/c), $E_{SC} < 1$ GeV and radiative Bhabhas (trailing cluster at $p < 1.6$ GeV/c, $E_{SC} \approx p$). To remove these backgrounds, a cut is made at $E_{SC} < 0.7$ GeV/c. In addition, the shower counter has a number of support ribs which are dead regions, thus degrading the energy measurement. Tracks which enter these regions are removed from consideration.

An additional handle on the identification of protons is gained from the dE/dx system. Figure 2 shows the reduced $\chi_{dE/dx}$ for candidate events that pass the above cuts. Reduced $\chi_{dE/dx}$ is defined as $\chi_{dE/dx}/DOF$. The vertical axis is for the $\overline{p}$ hypothesis, and the horizontal refers to the p hypothesis. The cluster near (0,0) contains true $p\overline{p}$ events, and the cluster near (5,5) is a mixture of event types such as radiative Bhabhas and dimuons. A cut is made on the combined χ^2, $(\chi_p^2 + \chi_{\overline{p}}^2)^{\frac{1}{2}} < 3$.

The weighted average momentum spectrum of the remaining candidate events is shown in Figure 3. By weighted average we mean that the original measurements on the positive and negative tracks (curvature and dip-angle) are averaged together and then combined to form a momentum. This spectrum in Figure 3 is fit to a gaussian plus a quadratic, with the centroid of the gaussian fit to the theoretic momentum of the protons, 1.586 GeV/c. The width and height are allowed to vary. From the fit, $N_{p\overline{p}} = 225 \pm 15 \pm 21$, where the first error is statistical and the second is the error on the fit.

B. $\psi' \to \Lambda\overline{\Lambda}$

The decays $\psi' \to \Lambda\overline{\Lambda}$ produce two back-to-back Λs, each with momentum 1.467 GeV/c. We only consider events where both Λs decay to the charged $p\pi$ final state. The final states of interest are thus, $\psi' \to \pi^+\pi^- p\overline{p}$, where the $p\pi^-$ and $\overline{p}\pi^+$ originate from well separated decay vertices. The decay kinematics are such that the proton (antiproton) is always the highest momentum positive (negative) track in the event.

1. $\Lambda\overline{\Lambda}$ selection

We select events with four and only four well reconstructed tracks with a zero net charge, and in the fiducial region covered by the drift chamber, $|\cos\theta| \leq 0.80$.

Events which pass these cuts are processed through a detached vertex finding algorithm, and subjected to a 5-C kinematic fit to $p\overline{p}\pi^+\pi^-$, with $M_{p\pi^-} = M_{\overline{p}\pi^+}$. The 84 events which pass this fit with a confidence level of more than 1%, and have $M_{p\pi} \leq 1.15$ GeV are shown

in Figure 4. Roughly extrapolating the two events in the region above 1.13 GeV/c to the area under the mass peak, we find that there are four background events in the plot. We conservatively assign this number a 100% error and determine $N_{\Lambda\overline{\Lambda}}$ to be $80 \pm 9 \pm 4$.

C. $\psi' \to \Sigma^0\overline{\Sigma}^0$

The Σ^0 hyperons from $\psi' \to \Sigma^0\overline{\Sigma}^0$ decay promptly via $\Sigma^0 \to \gamma\Lambda$. We consider only those decays where the daughter Λs decay via the charged $p\pi$ mode. The experimental signature is thus, $\psi' \to p\overline{p}\pi^+\pi^-\gamma\gamma$, where the $p\pi^-$ and $\overline{p}\pi^+$ originate from Λ hyperons with well separated decay vertices. In addition, there are two photons in the energy range $27 \leq E_\gamma \leq 202$ MeV. As in the case for $\psi' \to \Lambda\overline{\Lambda}$, the proton (antiproton) is always the highest momentum positive (negative) track in the final state.

1. $\Sigma^0\overline{\Sigma}^0$ Selection

We extract $\psi' \to \Sigma^0\overline{\Sigma}^0$ event candidates ($\Sigma^0 \to \Lambda\gamma$, $\Lambda \to p\pi^-$) using the same selection criteria as used for the $\Lambda\overline{\Lambda}$ mode with the additional requirement that there be two or more isolated clusters in the BSC with energy greater than 60 MeV, and within region $|\cos\theta| \leq 0.75$. By *"isolated"* we mean more than 12.8° ($\cos\theta_{\rm isol} < .975$) away from each of the charged tracks.

Both $p\pi$ pairs in the surviving events are processed through a displaced vertex-finding algorithm and the event is then subjected to a five-constraint kinematic to the hypothesis $\psi' \to p\overline{p}\pi^+\pi^-\gamma\gamma$, with the beam constraint $M_{p\pi^-\gamma} = M_{\overline{p}\pi^+\gamma}$. Here the highest momentum positive (negative) track is classified as the proton (antiproton). For events with more than two γ candidates, the fit is applied for each possible combination.

Events which pass the kinematic fit with a confidence level greater than 1%, and $M_{p\pi\gamma} < 1.3$ GeV/c^2 are shown in Figure 5. We fit this spectrum to a single gaussian plus a linear background with the peak fixed to the mass of the Σ^0, 1.192 GeV/c^2. From the fit, $N_{\Sigma^0\overline{\Sigma}^0} = 8 \pm 3 \pm 2$. The first error is statistical, and the second is the error on the fit.

D. $\psi' \to \Xi^-\overline{\Xi}^+$

The Ξ^- hyperon from $\psi' \to \Xi^-\overline{\Xi}^+$ decays via $\Xi^- \to \pi^-\Lambda$. We consider only those decays where the daughter Λs decay via the charged $p\pi$ mode. The experimental signature is thus $\psi' \to p\overline{p}\pi^+\pi^-\pi^+\pi^-$ where one each of the $p\pi^-$ and $\overline{p}\pi^+$ combinations originate from Λ hyperons with well separated decay vertices. As in the case for $\psi' \to \Lambda\overline{\Lambda}$ and $\Sigma^0\overline{\Sigma}^0$, the proton (antiproton) is always the highest momentum positive (negative) track in the final state.

1. $\Xi^-\overline{\Xi}^+$ *Selection*

We select events with six and only six well reconstructed tracks with zero net charge, and in the fiducial region covered by the drift chamber, $|\cos\theta| \leq 0.80$.

Each of the four possible $p\pi^-$ and $\overline{p}\pi^+$ combinations are sent through a displaced vertex-finding algorithm and subsequently subjected to a five-constraint kinematic fit to the hypothesis $\psi' \to p\overline{p}\pi^+\pi^-\pi^+\pi^-$, with the beam constraint $M_{p\pi^-\pi^-} = M_{\overline{p}\pi^+\pi^+}$. Due to kinematics, the highest momentum positive (negative) track is identified as the proton (antiproton).

Events which pass the fit with a confidence level greater than 1% are examined further. We additionally require that the $p\pi$ combinations have a mass within 10 MeV/c^2 of the Λ and that the mass of the $\Lambda\overline{\Lambda}$ candidate is more than 20 MeV/c^2 away from the J/ψ in order to reduce background from the cascade decay $\psi' \to J/\psi\pi\pi$, $J\psi \to \Lambda\overline{\Lambda}$.

The $M_{p\pi^-\pi^-}$ spectrum of events which remain after the above cuts is plotted in Figure 6. There are 12 ± 3.4 events in the Ξ^- peak. Averaging the five events outside the peak region over the entire plot and multiplying by the width of the signal gives 0.15 background events. A conservative error of 100 percent is applied, giving $12 \pm 3.4 \pm 0.2$ $\psi' \to \Xi^-\overline{Xi}^+$ events detected.

IV. BARYON DECUPLET

A. $\psi' \to \Delta^{++}\overline{\Delta}^{--}$

The decay $\psi' \to \Delta^{++}\overline{\Delta}^{--}$ produces two back-to-back Δ^{++} and $\overline{\Delta}^{--}$. As the Δ^{++} is a broad (111 MeV/c^2) resonance, the primary hyperons do not have well-defined momenta, in contrast to the octet cases above. We select events where both Δ^{++} and $\overline{\Delta}^{--}$ decay to $p\pi$ [$\mathcal{B}(\Delta^{++} \to p\pi) > 99\%$]. The final state is $\psi' \to p\overline{p}\pi^+\pi^-$.

1. Event Selection

We select events with four and only four well reconstructed tracks with a zero net charge, and in the fiducial region covered by the drift chamber, $|\cos\theta| \leq 0.80$. The surviving events are processed through a four-constraint kinematic fit to the hypothesis $\psi' \to p\overline{p}\pi^+\pi^-$. Events which pass with a confidence level greater than 1% are examined further.

Figure 8 shows the invariant mass distribution of the $p\overline{p}$ pair in events which pass the fit. There is a clear peak in the J/ψ mass region coming from the cascade decay $\psi' \to J/\psi\pi^+\pi^-$, $J/\psi \to p\overline{p}$; we remove this by making a 60 MeV/c^2 cut around the J/ψ. Figure 7 shows the invariant mass distribution for $p\pi^-$ containing a peak at the Λ mass. We remove the $\Lambda\overline{\Lambda}$ background by requiring the $p\pi^-$ and $\overline{p}\pi^+$ masses to be greater than 1.15 GeV/c^2.

Events which pass all above cuts are fit to a spin-1 Breit-Wigner plus a 4-body phase-space background histogram. The width and centroid of the signal spectrum are fixed to the PDG [6] values. Figure 9 shows the output of the fit; there are 849 total events in the plot. The fit parameter varied is the relative proportions of the phase space background and the Breit-Wigner signal to the total number of events in the plot. $N_{\Delta^{++}\overline{\Delta}^{--}}$ is $157 \pm 13 \pm 34$, where the first error is statistical and the second is the systematic error from the fitting procedure.

B. $\psi' \to \Sigma^+(1385)\overline{\Sigma}^-(1385)$

The hyperons from $\psi' \to \Sigma^{*+}\overline{\Sigma}^{*-}$ decay via $\Sigma^+(1385) \to \Lambda\pi^+$ 88% of the time. We consider only those decays where the daughter Λs decay via the charged $p\pi$ mode. The experimental signature is $\psi' \to p\overline{p}2(\pi^+\pi^-)$ where one each of the $p\pi^-$ ($\overline{p}\pi^+$) combinations are consistent with being from the decay of a Λ ($\overline{\Lambda}$).

1. $\Sigma^+(1385)\overline{\Sigma}^+(1385)$ *Selection*

We select events with six and only six well reconstructed tracks with a zero net charge, and in the fiducial region covered by the drift chamber, $|\cos\theta| \leq 0.80$. We kinematically fit the 36 possible charge combinations of $(+-)-(-+)+$, running the $(+-)/(-+)$ candidates through a displaced vertex finding algorithm, to $p\pi^-\pi^-\overline{p}\pi^+\pi^+$ with no other constraints.

Figure 10 shows that the fit mass of the daughter Λs from the decay of the primary Σ^{*+} is well defined and centered at the Λ mass. We make a loose cut of 15 MeV/c^2 on the Λ and $\overline{\Lambda}$ resonances.

As shown in Figure 11, the mass recoiling against the orphan $\pi^+\pi^-$ pair is dominated by a peak at the J/ψ mass, indicating contamination of $\psi' \to \pi^+\pi^- J/\psi$, $J/\psi \to \Lambda\overline{\Lambda}$. We therefore remove events with a $\pi^+\pi^-$ recoil mass within 30 MeV/c^2 of the J/ψ.

To determine $N_{\Sigma^{*+}\overline{\Sigma}^{*-}}$, we constrain the $\overline{p}\pi^+\pi^-$ combination ($\overline{\Sigma}^{*-}$ candidate) to be within 107.4 MeV/c^2 ($3 \times \Gamma$) of the nominal PDG value in order to enhance the Σ^{*+} signal. Events which pass the above cuts are fit to a Breit-Wigner with a constant background, with the mass and width fixed to the PDG values ($M = 1382.8$ GeV/c^2, $\Gamma = 35.8$ MeV/c^2). This fit is shown in Figure 12; from the fit, $N_{\Sigma^{*+}\overline{\Sigma}^{*-}} = 13.8 \pm 3.7 \pm 2.7$.

C. $\psi' \to \Xi^0(1530)\overline{\Xi}^0(1530)$

In the decay $\psi' \to \Xi^*\overline{\Xi}^*$, the Ξ^*s are produced back to back in the ψ' rest frame. The dominant decay mode of Ξ^* baryons is $\Xi^* \to \Xi^-\pi^+$, with a branching fraction of 0.66. [6]

The Ξ^- decays as in Section III D to $\Lambda\pi^-$, and the Λ decays to $p\pi^-$.

1. Event Selection

We select events with eight and only eight well reconstructed tracks with zero net charge, and in the fiducial region covered by the drift chamber, $|\cos\theta| \leq 0.80$. Remaining events are subjected to a 4-constraint kinematic fit to the hypothesis $\psi' \to p\pi^-\overline{p}\pi^+\pi^+\pi^-\pi^+\pi^-$. $p\pi$ candidates for Λs are sent through a displaced vertex finder. Events which pass the fit with a P_{fit} greater than 0.01 are examined further.

As the dominant decay mode in Ξ^* decays includes a Λ in the decay chain, a loose cut is placed on the $p\pi^-$ mass ($|M_{p\pi^-} - M_\Lambda| < 20$ MeV/c^2) to enhance the signal fraction (Figure 13). Due to π combinatorics, each event that passes the kinematic fit is counted four times in this plot.

Similarly, as there is a Ξ^- in the decay chain, a cut is made on the $p\pi^-\pi^-$ invariant mass; $M_{p\pi^-\pi^-}$ is required to be less than 20 MeV/c^2, as shown in Figure 14. Due to π combinatorics, each event is counted twice in this plot.

All events which remain after the above cuts are graphed in Figure 15 with $M_{\overline{p}\pi^+\pi^+\pi^-}$ on the vertical axis and $M_{p\pi^-\pi^-\pi^+}$ on the horizontal. The signal region is shown as a circle at (1.531,1.531). No events fall within the signal region defined as a 50 MeV/c^2 radius from the central value. We set an upper limit of 2.3 events at 90% CL for $N_{\Xi^*\overline{\Xi}^*}$.

D. $\psi' \to \Omega^-\overline{\Omega}^+$

The dominant Ω^- decay chain is $\Omega^- \to \Lambda K^-$, $\Lambda \to p\pi^-$ with a total branching fraction of 43% [6]. We look for $\psi' \to \Omega^-\overline{\Omega}^+$ events with the topology $\psi' \to p\overline{p}\pi^+\pi^-K^+K^-$, i.e. six charged tracks where the $p\pi^-$ and $\overline{p}\pi^+$ are consistent with being from the decay of a Λ or $\overline{\Lambda}$.

1. Event Selection

We select events with six charged tracks in the polar angle region $|\cos\theta| \leq 0.8$ and with zero net charge. The remaining events are subjected to a 7-constraint kinematic fit to the hypothesis $\psi' \to \Lambda\overline{\Lambda}K^+K^-$, $M_{\Lambda K^-} = M_{\overline{\Lambda}K^+}$. The fit is applied for each of the 36 particle assignment possibilities. Only the assignment with the best P_{fit} is considered.

Figure 16 shows the ΛK^- mass distribution for the selected events, where the solid line histogram is data and the crosses are from Monte Carlo, normalized to three events. There are no candidates within three sigma of the nominal Ω^- peak, thus an upper limit of 2.3 is quoted at eht 90 percent confidence level.

V. DETERMINATION OF THE TOTAL NUMBER OF ψ' EVENTS

We determine the number of ψ' events in our data samples from the observed number of cascade decays of the type $\psi' \to \pi^+\pi^- J/\psi$; $J/\psi \to \mu^+\mu^-$. The analysis for this is documented in reference [8].

The total number of ψ' events is determined to be $3.79 \pm 0.35 \times 10^6$ where the error is dominated by the error on the $\psi' \to \pi^+\pi^- J/\psi$ branching fraction.

VI. ACCEPTANCE AND EFFICIENCY

A. $p\overline{p}$

We determine the acceptance for $\psi' \to p\overline{p}$ events from a sample of Monte Carlo simulated events. Events were generated with a distribution of

$$\frac{dN}{d\cos\theta} \propto 1 + \alpha\cos^2\theta$$

with $\alpha = 0.61 \pm 0.23$. This proportionality constant was measured in the $J/\psi \to p\overline{p}$ system by [7]. Out of 20000 events generated, 14857 events survive these cuts, yielding a general

acceptance of 0.743. The collinearity cut is also purely geometric, and has an efficiency of 0.999.

As the BES Monte Carlo is of limited usefulness for simulating detailed hadronic interactions, cuts which are affected by such must be corrected for by the examination of real data. Fortunately, there is a subset of events which will allow the effects of these cuts to be determined. A clean sample of $p\overline{p}$ pairs was aquired from the analysis of $\Delta^{++}\overline{\Delta}^{--}$ in Section IV A. Instead of discarding the J/ψ contamination shown in Figure 8, these events are used to examine the E_{SC}, muon Id and dE/dx cuts.

This study is summarized in Table I. Values which could be determined via both methods were averaged, and the spread in the values used as a systematic error. Systematic error in other columns was determined from both Monte Carlo statistics and variation of cuts. The weighted product of the "Avg" column in Table I, 0.227 ± 0.032 is the efficiency in the $p\overline{p}$ mode.

B. $\Lambda\overline{\Lambda}$, $\Sigma^0\overline{\Sigma}^0$, and $\Xi^-\overline{\Xi}^+$

We determine the efficiency for the hyperon-pair channels completely from Monte Carlo simulated events. Here we generated 20000 events in each mode with a $1 + \alpha_d \cos\theta$ distribution, $\alpha_d = 0.67 \pm 0.21$, 0.22, 0.5 ± 0.5 for $\Lambda\overline{\Lambda}$, $\Sigma^0\overline{\Sigma}^0$, and $\Xi^-\overline{\Xi}^+$ respectively. α_d was not varied for the $\Sigma^0\overline{\Sigma}^0$ mode as the statistical error was large.

The resulting efficiencies are summarized in Table III. The systematic error reported is a combination of Monte Carlo statistics and variation of α_d.

In the three hyperon modes we require two Λ's that decay to charged $p\pi$ final states, which has a branching fraction $\mathcal{B}(\Lambda \to p\pi^-) = 0.639 \pm 0.005$; the other decay modes that are required are very nearly unity, namely $\mathcal{B}(\Sigma \to \Lambda\gamma) = 1.0$ and $\mathcal{B}(\Xi^- \to \Lambda\pi^-) = 0.999$ [6]. The acceptance for each channel is the $\Lambda \to p\pi^-$ branching fraction squared: 0.41 ± 0.01.

C. Acceptance and Efficiency of the Decuplet Pairs

We determine the efficiency for the decuplet hyperon-pair channels completely from Monte Carlo simulated events. Here we generated 20000 events in each mode with a $1 + \alpha_d \cos\theta$ distribution, α_d varying between 0 and 1 for $\Delta^{++}\overline{\Delta^{--}}$, and constant at 0.6 for the other modes. The resulting acceptances are summarized in Table III, where the systematic error reported is a combination of Monte Carlo statistics and variation of α_d.

The branching fraction for $\Delta^{++} \to \pi^+\pi^+$ is greater than 0.99 [6], thus the acceptance used is 1.00 ± 0.01. $\Sigma^{*+}\overline{\Sigma}^{*-}$ contains two Λs going to $p\pi$ (0.639^2) and two Σ^*s decaying to $\Lambda\pi$ (0.88^2), for a total acceptance of 0.316 ± 0.011. $\Xi^{*0}\overline{\Xi}^{*0}$ has 3 components to the acceptance: $\Xi^{*0} -> \Xi^-\pi^+$ (0.650^2), $\Xi^- \to \Lambda\pi^-$ (0.999^2) $\Lambda \to p\pi^-$ (0.639^2), with a total acceptance of 0.172 ± 0.001. $\Omega^-\overline{\Omega}^+$ has only two components in the acceptance, $\Omega^- \to \Lambda K^-$ (0.678^2) and $\Lambda \to p\pi^-$ (0.639^2), with a total acceptance of 0.187 ± 0.001.

VII. RESULTS

The branching fractions are determined from the expression

$$\mathcal{B}(\psi' \to X) = \frac{N_X}{N_{\psi'}\epsilon\eta},$$

where N_X, $N_{\psi'}$, and ϵ are the numbers of selected events, the total number ψ's, and the combined acceptance and efficiency, respectively. The values of these quantities and the resulting branching fraction for each decay mode are listed in Table III.

We compare our results for the branching fractions to previous limits and results in Figure 17. Our measured value for the $\mathcal{B}(\psi' \to p\overline{p})$ is about one standard deviation higher than the previous DASP measurements, which was based on 4 events [3] and a Mark I measurement with similar statistics [4]. The results for $\Lambda\overline{\Lambda}$ and $\Xi^-\overline{\Xi}^-$ are within the PDG upper limit values. There are no previous experimental results for $\Sigma^0\overline{\Sigma}^0$ or any of the decuplet modes.

In Fig. 18, we plot the reduced branching fractions derived from our measurements. The results show a trend to smaller values for the higher massses, similar to that seen for the J/ψ and is only marginally consistent with expectations from flavor-$SU(3)$ symmetry. Higher precision measurements *both* for the J/ψ and ψ' would clarify this issue.

Our measured ψ' branching fractions agree with expectations derived from the application of the 14% rule to the PDG values for the corresponding J/ψ decays, shown in Table IV.

A. Results

The values of the resulting branching fraction for each decay mode are listed in Table III.

The N_{evt} entries for the modes $\Sigma^0\overline{\Sigma}^0$, $\Delta^{++}\overline{\Delta}^{--}$, and $\Sigma^{*-}\overline{\Sigma}^{*-}$ are already background subtracted. We plot our results for the branching fractions in Fig. 19, There are no previous experimental results for those modes. Currently we only give upper limits for $\Xi^{*0}\overline{\Xi}^{*0}$ and $\Omega^-\overline{\Omega}^+$, we need more ψ' data to study these two modes.

Our measured ψ' branching fractions agree with expectations derived from the application of the 14% rule to the PDG values for the corresponding J/ψ decays for $\Delta^{++}\overline{\Delta}^{--}$ and $\Sigma^{*-}\overline{\Sigma}^{*+}$, no entry in J/ψ to $\Xi^0(1530)\overline{\Xi}^0(1530)$ and $\Omega^-\overline{\Omega}^+$.

VIII. CONCLUSIONS

We report preliminary measurements of the branching fractions for $\psi' \to p\overline{p}$, $\Lambda\overline{\Lambda}$, $\Sigma^0\overline{\Sigma}^0$, and $\Xi^-\overline{\Xi}^+$. The measured branching fractions agree with expectations based on an application of the 14% rule to the corresponding J/ψ decays. The reduced branching fractions decrease with increasing baryon masses, showing some deviation from expectations based on flavor-$SU(3)$ symmetry.

IX. ACKNOWLEDGEMENTS

We thank the staffs of the BEPC accelerator and the IHEP computing center for their assistance.

REFERENCES

† Work supported in part by the National Natural Science Foundation of China under Contract No. 19290400 and the Chinese Academy of Sciences under contract No. KJ85 (IHEP); by the Department of Energy under Contract Nos. DE-FG02-91ER40676 (Boston University), DE-FG03-92ER40701 (Caltech), DE-FG03-93ER40788 (Colorado State University), DE-AC02-76ER03069 (MIT), DE-AC03-76SF00515 (SLAC), DE-FG03-91ER40679 (UC Irvine), DE-FG03-94ER40833 (U Hawaii), DE-FG05-92ER40736 (UT Dallas), DE-AC35-89ER40486 (SSC Lab); by the U.S. National Science Foundation, Grant No. PHY9203212 (University of Washington); and by the Texas National Research Laboratory Commission under Contract Nos. RGFY91B5, RGFY92B5 (Colorado State), and RCFY93-316H (UT Dallas).

[1] α_s is calculated using the method given by the Particle Data Group [6], with $\Lambda^{(4)} = 280$ MeV. The ratio $\alpha_s(\psi')/\alpha_s(J\psi)$ is calculated to be about 0.8 by this method.

[2] M.E.B. Franklin et al. (Mark II Collaboration), Phys. Rev. Lett. **51**, 963 (1983).

[3] R. Brandelick et al. (DASP Collaboration), Z. Phys. **C1**, 233 (1979).

[4] G. Feldman and M. Perl (Mark I Collaboration), Phys. Rep. **C33**, 285 (1977).

[5] J.Z. Bai et al. (BES Collaboration), Nucl. Inst. and Meth. **A344**, 319 (1994).

[6] R. M. Barnett *et al.* (Particle Data Group), *Review of Particle Physics,* Phys. Rev. **D54**, 1 (1996).

[7] M.W. Eaton et al. (Mark II Collaboration), Phys. Rev. **D29**, 804 (1984).

[8] J.Z. Bai et al. (BES Collaboration), Phys. Rev. **D58**:092006 (1998).

FIGURES



FIG. 1. Shower counter energy *vs.* momentum for positively charged tracks.



FIG. 2. dE/dx reduced χ for diproton candidates. Vertical axis is for $\overline{p}$ candidates, horizontal is for p candidates. A cut is made on the combined χ^2, $(\chi_p^2 + \chi_{\overline{p}}^2)^{\frac{1}{2}} < 3$.



FIG. 3. Weighted average momentum of $p\overline{p}$ pairs, fit to a gaussian plus a quadratic.



FIG. 4. $p\pi$ invariant mass from a kinematic fit to $\psi' \to p\overline{p}\pi^+\pi^-$, $M_{p\pi^-} = M_{\overline{p}\pi^+}$.



FIG. 5. $p\pi\gamma$ invariant mass from a kinematic fit to $\psi' \to p\overline{p}\pi^+\pi^-\gamma\gamma$, $M_{p\pi^-\gamma} = M_{\overline{p}\pi^+\gamma}$. Events with $p\pi\gamma$ masses below 1.3 GeV/c^2 are fit to a gaussian signal plus a linear background.



FIG. 6. $p\pi^-\pi^-$ masses from a kinematic fit to $\psi' \to p\overline{p}\pi^+\pi^-\pi^+\pi^-$, $M_{p\pi^-\pi^-} = M_{\overline{p}\pi^+\pi^+}$. There are $12 \pm 3.4 \pm 0.2$ events in the Ξ^- peak.



FIG. 7. $p\pi^-$ masses in the Δ^{++} analysis. A cut is made at $M_{p\pi^-} > 1.14$ GeV/c^2 to remove $\Lambda\overline{\Lambda}$ background.



FIG. 8. $\pi^+\pi^-$ recoil masses in the Δ^{++} analysis. A cut is made at $|M_{p\overline{p}} - M_{J\psi}| > 60$ MeV/c^2 to remove J/ψ contamination.



FIG. 9. $p\pi^+$ masses from a kinematic fit to $\psi' \to p\overline{p}\pi^+ pi^-$. Data is fit to a spin-1 Breit-Wigner plus a 4-body phase-space background. $N_{\Delta^{++}\overline{\Delta}^{--}} = 157 \pm 13 \pm 34$.



FIG. 10. $M_{p\pi^-}$ from a fit to $\psi' \to (p\pi^-)\pi^-(\overline{p}\pi^+)\pi^+$.



FIG. 11. $\pi^+\pi^-$ recoil mass from a kinematic fit to $\psi' \to \Lambda\overline{\Lambda}\pi^+\pi^-$, showing large J/ψ contamination.



FIG. 12. $M_{p\pi^-\pi^-}$ from a fit to $\psi' \to (p\pi^-)\pi^-(\overline{p}\pi^+)\pi^+$.



FIG. 13. $p\pi^-$ masses from a kinematic fit to $\psi' \to p\overline{p}\pi^+\pi^-\pi^+\pi^-\pi^+\pi^-$.



FIG. 14. $p\pi^-\pi^-$ masses from a kinematic fit to $\psi' \to p\overline{p}\pi^+\pi^-\pi^+\pi^-\pi^+\pi^-$.



FIG. 15. $M_{\overline{p}\pi^+\pi^+\pi^-}$ vs $M_{p\pi^-\pi^-\pi^+}$ from a kinematic fit to $\psi' \to p\overline{p}\pi^+\pi^-\pi^+\pi^-\pi^+\pi^-$. The circle denotes the signal region.



FIG. 16. ΛK^- mass distribution from a kinematic fit to $\psi' \to \Lambda\overline{\Lambda}K^+K^-$. Histogram is data, crosses are Monte Carlo.



FIG. 17. Comparison of measured branching fractions (circles) with previous measurements (triangles).



FIG. 18. The reduced branching fractions $|M_i|^2$ for $\psi' \to B_i\overline{B_i}$ decays.

FIG. 19. The measured branching fractions for $\psi' \to B_i\overline{B_i}$ decays,$B_i = \Delta^{++}$, $\Sigma^{*\,+}$, $\Xi^{0\,*}$ and Ω^-.

FIG. 20. The reduced branching fractions $|M_i|^2$ for $\psi' \to B_i\overline{B_i}$ decuplet decays.

TABLES

Cut	MC	$\epsilon_{\rm MC}$	J/ψ	$\epsilon_{J/\psi}$	Avg	Syst
None	20000					
General	14857	.743	847		.743	.006
Muon ID	10335	.696	662	.782	.739	.043
BSC Geom	7942	.768	418	.631	.700	.069
$E_{SC} < 0.7$			404	.610	.610	
$\|XSP\| < 3$			391	.968	.968	.047
Collinearity	7937	.999			.999	.011

TABLE I. Relative efficiencies for cuts in the mode $\psi' \to p\overline{p}$, as modeled by Monte Carlo and J/ψ data. Avg column denotes efficiency used in the analysis.

Channel	Decay mode	Topology	Fraction
$p\overline{p}$		$p\overline{p}$	1.0
$\Lambda\overline{\Lambda}$	$\Lambda \to p\pi^-$	$p\overline{p}\pi^+\pi^-$	0.41
$\Sigma^0\overline{\Sigma}^0$	$\Sigma^0 \to \Lambda\gamma, \Lambda \to p\pi^-$	$p\overline{p}\pi^+\pi^-\gamma\gamma$	0.41
$\Xi^-\overline{\Xi}^+$	$\Xi^- \to \Lambda\pi^+, \Lambda \to p\pi^-$	$p\overline{p}2(\pi^+\pi^-)$	0.41
$\Delta^{++}\overline{\Delta}^{--}$	$\Delta^{++} \to p\pi^+, \overline{\Delta}^{--} \to \overline{p}\pi^-$	$p\overline{p}\pi^+\pi^-$	>0.98
$\Sigma^{*+}\overline{\Sigma}^{*-}$	$\Sigma^* \to \Lambda\pi, \Lambda \to p\pi^-$	$p\overline{p}2(\pi^+\pi^-)$	0.64
$\Xi^{*0}\overline{\Xi}^{*0}$	$\Xi^{*0} \to \Xi^-\pi^+, \Xi^- \to \Lambda\pi^-$	$p\overline{p}3(\pi^+\pi^-)$	0.41
$\Omega^-\overline{\Omega}^+$	$\Omega^- \to \Lambda K^-, \Lambda \to p\pi^-$	$p\overline{p}\pi^+\pi^-K^+K^-$	0.19

TABLE II. Decay mode, final topology, and decay fraction for baryonic initial states.

mode	N_{evt}	efficiency	B.F. Acceptance	$\mathcal{B}(\times 10^{-5})$
$p\overline{p}$	$225 \pm 15 \pm 21$	0.227 ± 0.032	1.00	$26.2 \pm 1.7 \pm 5.5$
$\Lambda\overline{\Lambda}$	$80 \pm 9 \pm 4$	0.27 ± 0.01	0.41 ± 0.01	$18.9 \pm 2.1 \pm 2.1$
$\Sigma^0\overline{\Sigma}^0$	$8 \pm 3 \pm 2$	0.043 ± 0.003	0.41 ± 0.01	$12 \pm 4 \pm 3$
$\Xi^-\overline{\Xi}^+$	$12 \pm 3.4 \pm 0.2$	0.078 ± 0.01	0.41 ± 0.01	$10 \pm 3 \pm 1$
$\Delta^{++}\overline{\Delta}^{--}$	$157 \pm 13 \pm 34$	0.31 ± 0.02	1.0 ± 0.01	$13.4 \pm 1.1 \pm 3.3$
$\Sigma^{*+}\overline{\Sigma}^{*-}$	$14 \pm 4 \pm 3$	0.104 ± 0.005	0.316 ± 0.011	$11 \pm 3 \pm 3$
$\Xi^{*0}\overline{\Xi}^{*0}$	< 2.3	0.041 ± 0.001	0.172 ± 0.001	< 8.5
$\Omega^-\overline{\Omega}^+$	< 2.3	0.042 ± 0.001	0.187 ± 0.001	< 7.7

TABLE III. Results for $\psi' \to B_i\overline{B_i}$.

Decay Mode	$\mathcal{B}$	$0.147 \times \mathcal{B}$
$J/\psi \to \quad p\overline{p}$	$(2.14 \pm 0.10) \times 10^{-3}$	$(3.15 \pm 0.15) \times 10^{-4}$
$J/\psi \to \quad \Lambda\overline{\Lambda}$	$(1.35 \pm 0.14) \times 10^{-3}$	$(1.98 \pm 0.21) \times 10^{-4}$
$J/\psi \to \quad \Sigma^0\overline{\Sigma}^0$	$(1.3 \pm 0.2) \times 10^{-3}$	$(1.9 \pm 0.3) \times 10^{-4}$
$J/\psi \to \quad \Xi^-\overline{\Xi}^+$	$(0.9 \pm 0.2) \times 10^{-3}$	$(1.3 \pm 0.3) \times 10^{-4}$.
$J/\psi \to \Delta^{++}\overline{\Delta}^{--}$	$(1.10 \pm 0.29) \times 10^{-3}$	$(1.62 \pm 0.5) \times 10^{-4}$
$J/\psi \to \quad \Sigma^-\overline{\Sigma}^+$	$(1.03 \pm 0.13) \times 10^{-3}$	$(1.40 \pm 0.24) \times 10^{-4}$

TABLE IV. Branching ratio comparisons between J/ψ and ψ'.