Study of the P-wave charmonium state χ_{cJ} in $\psi(2S)$ decays

J. Z. Bai[1], J. G. Bian[1], I. Blum[11], Z. W. Chai[1], G. P. Chen[1], H. F. Chen[10], J. Chen[3], J. C. Chen[1], Y. Chen[1], Y. B. Chen[1], Y. Q. Chen[1], B. S. Cheng[1] X. Z. Cui[1], H. L. Ding[1], L. Y. Ding[1], L. Y. Dong[1], Z. Z. Du[1], W. Dunwoodie[7], S. Feng[1], C. S. Gao[1], M. L. Gao[1], S. Q. Gao[1], P. Gratton[11], J. H. Gu[1], S. D. Gu[1], W. X. Gu[1], Y. F. Gu[1], Y. N. Guo[1], S. W. Han[1], Y. Han[1], F. A. Harris[8], J. He[1], J. T. He[1], M. He[5], D. G. Hitlin[2], G. Y. Hu[1], H. M. Hu[1], J. L. Hu[1], Q. H. Hu[1], T. Hu[1], X. Q. Hu[1], J. D. Huang[1], Y. Z. Huang[1], J. M. Izen[11], C. H. Jiang[1], Y. Jin[1], Z. J. Ke[1], M. H. Kelsey[2], B. K. Kim[11], D. Kong[8], Y. F. Lai[1], P. F. Lang[1], A. Lankford[9], C. G. Li[1], D. Li[1], H. B. Li[1], J. Li[1], P. Q. Li[1], R. B. Li[1], W. Li[1], W. D. Li[1], W. G. Li[1], X. H. Li[1], X. N. Li[1], H. M. Liu[1], J. Liu[1], J. H. Liu[1], R. G. Liu[1], Y. Liu[1], X. C. Lou[11], B. Lowery[11], F. Lu[1], J. G. Lu[1], J. Y. Lu[1], L. C. Lu[1], C. H. Luo[1], A .M .Ma[1], E. C. Ma[1], J. M. Ma[1], R. Malchow[3], H. S. Mao[1], Z. P. Mao[1], X. C. Meng[1], J. Nie[1], S. L. Olsen[8], J. Oyang[2], D. Paluselli[8], L. J. Pan[8], J. Panetta[2], F. Porter[2], N. D. Qi[1], X. R. Qi[1], C. D. Qian[6], J. F. Qiu[1], Y. H. Qu[1], Y. K. Que[1], G. Rong[1], M. Schernau[9], Y. Y. Shao[1], B. W. Shen[1], D. L. Shen[1], H. Shen[1], X. Y. Shen[1], H. Y. Sheng[1], H. Z. Shi[1], X. F. Song[1], J. Standifird[11], F. Sun[1], H. S. Sun[1], S. Q. Tang[1], W. Toki[3], G. L. Tong[1], F. Wang[1], L. S. Wang[1], L. Z. Wang[1], M. Wang[1], Meng Wang[1], P. Wang[1], P. L. Wang[1], S. M. Wang[1], T. J. Wang[1†], Y. Y. Wang[1], M. Weaver[2], C. L. Wei[1], Y. G. Wu[1], D. M. Xi[1], X. M. Xia[1], P. P. Xie[1], Y. Xie[1], Y. H. Xie[1], W. J. Xiong[1], C. C. Xu[1], G. F. Xu[1], S. T. Xue[1], J. Yan[1], W. G. Yan[1], C. M. Yang[1], C. Y. Yang[1], J. Yang[1], W. Yang[3], X. F. Yang[1], M. H. Ye[1], S. W. Ye[10], Y. X. Ye[10], K. Yi [1], C. S. Yu[1], C. X. Yu[1], Y. H. Yu[4], Z. Q. Yu[1], Z. T. Yu[1], C. Z. Yuan[1], Y. Yuan[1], B. Y. Zhang[1], C. C. Zhang[1], D. H. Zhang[1], Dehong Zhang[1], H. L. Zhang[1], J. Zhang[1], J. L. Zhang[1], J. W. Zhang[1], L. S. Zhang[1], Q. J. Zhang[1], S. Q. Zhang[1], X. Y. Zhang[5], Y. Zhang[1], Y. Y. Zhang[1], D. X. Zhao[1], H. W. Zhao[1], J. W. Zhao[1], M. Zhao[1], W. R. Zhao[1], Z. G. Zhao[1], J. P. Zheng[1], L. S. Zheng[1], Z. P. Zheng[1], G. P. Zhou[1], H. S. Zhou[1], L. Zhou[1], Q. M. Zhu[1], Y. C. Zhu[1], Y. S. Zhu[1], B. A. Zhuang[1]

(BES Collaboration)

[1] *Institute of High Energy Physics, Beijing 100039, People's Republic of China*

[2] *California Institute of Technology, Pasadena, California 91125*

[3] *Colorado State University, Fort Collins, Colorado 80523*

[4] *Hangzhou University, Hangzhou 310028, People's Republic of China*

[5] *Shandong University, Jinan 250100, People's Republic of China*

[6] *Shanghai Jiaotong University, Shanghai 200030, People's Republic of China*

[7] *Stanford Linear Accelerator Center, Stanford, California 94309*

[8] *University of Hawaii, Honolulu, Hawaii 96822*

[9] *University of California at Irvine, Irvine, California 92717*

[10] *University of Science and Technology of China, Hefei 230026, People's Republic of China*

[11] *University of Texas at Dallas, Richardson, Texas 75083-0688*

(July 3, 1998)

Abstract

The processes $\psi(2S) \to \gamma\pi^+\pi^-$, γK^+K^- and $\gamma p\bar{p}$ have been studied using a sample of 3.7×10^6 produced $\psi(2S)$ decays. We determine the total width of the χ_{c0} to be $\Gamma^{tot}_{\chi_{c0}} = 14.3 \pm 2.0 \pm 3.0$ MeV. We present the first measurement of the branching fraction $B(\chi_{c0} \to p\bar{p}) = (16.3 \pm 4.4 \pm 5.4) \times 10^{-5}$, where the first error is statistical and the second one systematic. Branching fractions of $\chi_{c0,2} \to \pi^+\pi^-$ and K^+K^- are also reported.

Typeset using REVTEX

The hadronic decay rates of the P-wave quarkonium states provide tests of perturbative QCD. Recently, a systematic approach to the treatment of the infrared ambiguities in the calculation of the production and decays of these states has been developed [1]. However, existing experimental information on the triplet P-wave $c\bar{c}$ states ($\chi_{c0,1,2}$), especially the $J = 0$ χ_c, is not adequate for testing the predictions of this new theory.

In particular, the total width of the χ_{c0} is a quantity of considerable interest. However, the two existing measurements have large errors and only marginal consistency [2]. Also of interest is the decay $\chi_{c0} \to p\bar{p}$, which is forbidden in the limit of massless helicity conservation [3]. Here the only existing measurement is an upper limit on the partial width that does not seriously constrain the theory [4]. Calculation of the branching fractions for other exclusive χ_{cJ} decays, such as $\chi_{cJ} \to \pi^+\pi^-$ [5], has revealed orders-of-magnitude discrepancies with the data reported by early experiments. For these reasons, measurements of these properties of the χ_{cJ} states with improved precision are very useful.

In this paper we report a measurement of $\Gamma^{tot}_{\chi_{c0}}$ determined from an analysis of exclusive $\psi(2S) \to \gamma\pi^+\pi^-$ and γK^+K^- decays seen in the Beijing Spectrometer (BES) at the Beijing Electron Positron Collider (BEPC). We also report a first measurement of the $\chi_{c0} \to p\bar{p}$ branching fraction, and improved precision on the branching fractions for $\chi_{c0,2} \to \pi^+\pi^-$ and K^+K^-.

The BES is a conventional solenoidal magnet detector that is described in detail in ref. [6]. A four-layer central drift chamber (CDC) surrounding the beampipe provides trigger information. A forty-layer main drift chamber (MDC), located radially outside the CDC, provides trajectory and energy loss (dE/dx) information for charged tracks over 85% of the total solid angle. The momentum resolution is $\sigma_p/p = 0.017\sqrt{1+p^2}$ (p in GeV/c), and the dE/dx resolution for hadron tracks is $\sim 11\%$. An array of 48 scintillation counters surrounding the MDC measures the time-of-flight (TOF) of charged tracks with a resolution of ~ 450 ps for hadrons. Radially outside the TOF system is a 12 radiation length, lead-gas barrel shower counter (BSC). This measures the energies of electrons and photons over $\sim 80\%$ of the total solid angle with an energy resolution of $\sigma_E/E = 22\%/\sqrt{E}$ (E in GeV). Outside of the solenoidal coil, which provides a 0.4 Tesla magnetic field over the tracking volume, is an iron flux return that is instrumented with three double layers of counters that identify muons of momentum greater than 0.5 GeV/c.

We study χ_c states produced by the reaction $e^+e^- \to \psi(2S) \to \gamma\chi_c$ in a data sample corresponding to a total of 3.7 million $\psi(2S)$ decays. For the $\Gamma^{tot}_{\chi_{c0}}$ determination reported here we use the paired pseudoscalar meson decay modes of $\chi_{c0,2} \to \pi^+\pi^-$ and K^+K^-. Using the particle identification capabilities of the detector and four-constraint kinematic fits, we can get relatively pure event samples. Moreover, since the decays of the χ_{c1} and the η_c' to $\pi^+\pi^-$ or K^+K^- are forbidden by parity conservation, the χ_{c0} and χ_{c2} signals in these channels are free of distortions due to possible contamination of these other states. The effects of cross-contamination between the $\pi^+\pi^-$ and K^+K^- event samples are estimated by Monte Carlo (MC) simulations and corrected accordingly. The total width of the χ_{c2} has been precisely measured to be $\Gamma^{tot}_{\chi_{c2}} = 2.00 \pm 0.18$ MeV [7], which is much narrower than our experimental resolution at $M_{\chi_{c2}}$ of 7.83 MeV. Thus, the strong $\chi_{c2} \to \pi^+\pi^-$ and K^+K^- signals in our data are used to provide a direct experimental determination of our resolution. We only rely on the MC simulation to determine how the resolution changes between $M_{\chi_{c2}}$ and $M_{\chi_{c0}}$.

We select $\psi(2S) \to \gamma\pi^+\pi^-$, γK^+K^- and $\gamma p\bar{p}$ by imposing the following selection criteria.

A cluster of deposited energy in the BSC is regarded as a photon candidate if:

1. the angle between the nearest charged track and the cluster in the $r\phi$ plane is greater than 15°;

2. the energy of the cluster is greater than 20 MeV and some energy is deposited in the first 6 radiation lengths of the counter; and

3. the angle determined from the cluster development in the BSC agrees with that determined from the relative position of the shower location and the interaction point to within 37°.

At least one and at most three photon candidates are allowed in an event. The candidate with the largest BSC energy is assumed to be the photon radiated from the $\psi(2S)$.

In addition, we require that the event has two oppositely signed charged tracks in the MDC that both have at least 13 good hits and are well fit to a three dimensional helix. Events with tracks where the dE/dx measured in the MDC and the shower properties in the BSC are consistent with electrons are rejected. For each track, the TOF and dE/dx measurements are used to assign probabilities that the particle is a pion, kaon and proton ($Prob_\pi, Prob_K, Prob_p$). We require both tracks to have $Prob_\pi > 0.01$ (for $\pi^+\pi^-$) or $Prob_K > 0.01$ (for K^+K^-) or $Prob_p > 0.05$ (for $p\bar{p}$). In addition we do four-constraint kinematic fits to the hypotheses $\psi(2S) \to \gamma\pi^+\pi^-$, $\psi(2S) \to \gamma K^+K^-$ and $\psi(2S) \to \gamma p\bar{p}$ and require the χ^2 probability of the fit P_{χ^2} to be greater than 0.01 for $\pi^+\pi^-$ or K^+K^- and greater than 0.05 for $p\bar{p}$.

There is some background from $J/\psi \to \mu^+\mu^-$, where the J/ψ is produced by cascade $\psi(2S)$ to J/ψ decays. To reduce this, we reject events where the response of the muon detection system is consistent with the two charged tracks being muons. The surviving $\mu^+\mu^-$ background events do not populate the $\pi^+\pi^-$ invariant mass distribution near the χ_{c0} or χ_{c2} masses. In the K^+K^- mass distribution, however, they populate the region in the lower mass side of χ_{c0}, and cause an abnormal distribution. For the $p\bar{p}$ sample, the $\mu^+\mu^-$ background level is significant. For this channel, in order to insure that both tracks are directed at the sensitive area of the muon detection system, we require $|\cos\theta_{\rm MDC}| < 0.65$ for both the p and the $\bar{p}$ track.

To distinguish $\gamma\pi^+\pi^-$ from γK^+K^-, we define

$$Prob_{all}^{P^+P^-} = Prob(\chi^2_{\rm all}, ndf_{\rm all}),$$

where $\chi^2_{\rm all} = \chi^2_{4C} + \chi^2_{TOF} + \chi^2_{dE/dx}$ and $ndf_{\rm all} = ndf_{4C} + ndf_{TOF} + ndf_{dE/dx}$ are the total χ^2 and the corresponding number of degrees of freedom of the χ^2 distribution. Here χ^2_{4C}, χ^2_{TOF} and $\chi^2_{dE/dx}$ correspond to the χ^2 values from the 4-constraint kinematic fit, the TOF measurements for the π or K hypothesis, and the dE/dx measurements for the π or K hypothesis, respectively, and ndf_{4C}, ndf_{TOF} and $ndf_{dE/dx}$ are the corresponding numbers of degrees of freedom. If $Prob_{all}^{\pi^+\pi^-} > Prob_{all}^{K^+K^-}$, the event is categorized as a $\gamma\pi^+\pi^-$ event, and if $Prob_{all}^{K^+K^-} > Prob_{all}^{\pi^+\pi^-}$, it is categorized as a γK^+K^- event.

After the imposition of all of the above-listed selection requirements, Figs. 1 and 2 show the $\pi^+\pi^-$ and K^+K^- invariant mass distributions, respectively. In these plots, the mass values corresponding to the χ_{c0} and χ_{c2} peaks are lower in the $\pi^+\pi^-$ channel and higher for K^+K^-, indicating the presence of some remaining cross contamination between the two samples, which must be accounted for in the determination of the χ_{c0} parameters. From

Fig. 2, it is apparent that the $\chi_{c2} \to K^+K^-$ sample is statistically limited. We therefore only use the $\chi_{c2} \to \pi^+\pi^-$ signal to calibrate the mass resolution. In Fig. 3, the $p\bar{p}$ invariant mass distribution, there is a clear χ_{c0} signal and evidence for the χ_{c1} and χ_{c2}.

We use Monte Carlo simulated data to determine the $\pi^+\pi^-$ and K^+K^- cross contamination probabilities, the detection efficiencies and the mass resolutions. We generate events assuming that the reaction $\psi(2S) \to \gamma\chi_{cJ}$ is a pure E1 transition. The decays $\chi_{cJ} \to$ pseudoscalar meson pairs have only one independent helicity amplitude and are thus unambiguous [8]. For $\chi_{cJ} \to p\bar{p}$ decays, there are no available experimental data on the helicity amplitudes, and we use an isotropic distribution. (Our $p\bar{p}$ event samples are too small to permit a helicity amplitude analysis.)

We subject the MC-generated events to the same selection process as is used for the data and determine the detection efficiencies for each mode. For the $p\bar{p}$ mode, the detection efficiencies are $\varepsilon_{\chi_{c0}} = 27.1\%$, $\varepsilon_{\chi_{c1}} = 30.3\%$ and $\varepsilon_{\chi_{c2}} = 27.6\%$, and mass resolutions at the χ_{c0}, χ_{c1} and χ_{c2} equal to 7.3, 6.8 and 6.7 MeV, respectively. For the $\pi^+\pi^-$ and K^+K^- modes, the simulation shows that the mass dependence of the detection efficiency is small and the mass resolution function is very nearly Gaussian. We compensate for the distortion of the mass spectra due to the the $\pi^+\pi^-$-K^+K^- cross contamination by calibrating the mass resolution derived from the MC simulation with the $\chi_{c2} \to \pi^+\pi^-$ line shape seen with the data. The efficiencies are $\varepsilon_{\chi_{c0}\to\pi^+\pi^-} = 36.9\%$, $\varepsilon_{\chi_{c2}\to\pi^+\pi^-} = 38.9\%$, $\varepsilon_{\chi_{c0}\to K^+K^-} = 32.8\%$ and $\varepsilon_{\chi_{c2}\to K^+K^-} = 34.9\%$, and the probability for χ_{c0} (χ_{c2}) $\to K^+K^-$ events to be categorized as $\pi^+\pi^-$ is 5.6% (6.0%), and that for χ_{c0} (χ_{c2}) $\to \pi^+\pi^-$ events to be selected as K^+K^- is 7.1% (7.4%).

The invariant mass distributions in Figs. 1, 2 and 3 are fit by using an unbinned maximum likelihood algorithm. For the $p\bar{p}$ channel, the invariant mass region between 3.20 and 3.64 GeV is fit with three Breit-Wigner resonances plus a linear background function. The Breit-Wigner resonance width for the χ_{c0} is fixed at 14.3 MeV, the value determined from an analysis of $\chi_{c0} \to \pi^+\pi^-$ decays described below; those for the χ_{c1} and χ_{c2} are fixed at the PDG values [7]. The resonances are smeared by gaussian functions with rms widths fixed at the MC-determined mass resolution values. The fit result, shown as the curve in Fig. 3, gives $15.2 \pm 4.1, 4.2 \pm 2.2$ and 4.7 ± 2.5 events for the χ_{c0}, χ_{c1} and χ_{c2} states, respectively.

For the $\pi^+\pi^-$ channel, we first fit the invariant mass region between 3.5 and 3.6 GeV with a Breit-Wigner resonance with $\Gamma_{\chi_{c2}}$ fixed at the PDG value of 2.00 MeV, smeared by a Gaussian resolution function with an rms width that is allowed to float. We also include a linear background function. The fit results in a χ_{c2} mass resolution of 7.83 ± 1.04 MeV, which is slightly higher than the MC result of 6.31 ± 0.11 MeV. We scale the MC value for the mass resolution at the χ_{c0} (8.12 ± 0.23 MeV) by the ratio of the fitted to MC results at the χ_{c2} and get a mass resolution at the χ_{c0} of 10.08 MeV.

We then fit the $\pi^+\pi^-$ mass spectrum between 3.2 and 3.6 GeV with two gaussian-smeared Breit-Wigners with resolutions fixed at 10.08 and 7.83 MeV, and a second order polynomial background function, and with $\Gamma_{\chi_{c2}}$ fixed at the PDG value of 2.00 MeV. The fit, shown as the curve in Fig. 1, gives $\Gamma_{\chi_{c0}} = 14.3 \pm 2.0$ MeV, where the error is statistical. The fitted numbers of $\chi_{cJ} \to \pi^+\pi^-$ events are $n^{obs}_{\chi_{c0}\to\pi^+\pi^-} = 720 \pm 32$ and $n^{obs}_{\chi_{c2}\to\pi^+\pi^-} = 185 \pm 16$, where the errors are statistical.

We fit the K^+K^- mass spectrum between 3.2 and 3.6 GeV to two Gaussian-smeared Breit-Wigner resonance functions plus a background function that includes the possibility

of distortions to the line shape due to $\mu^+\mu^- X$ background events. (Because this mode is not used for width determination, a high precision knowledge of the mass resolution is not an issue.) The width of χ_{c2} is fixed. The resulting fit, shown in Fig. 2, gives $n^{obs}_{\chi_{c0}\to K^+K^-} = 774 \pm 38$ and $n^{obs}_{\chi_{c2}\to K^+K^-} = 115 \pm 13$, where the errors are statistical.

Errors in the determination of $\Gamma_{\chi_{c0}}$ are caused by the uncertainty in $\Gamma_{\chi_{c2}}$, the determination of the mass resolution, the shape of the background, the mass dependence of the efficiency correction, and the choice of experimental cuts. We add the estimated errors from these sources in quadrature and get a total relative systematic error on $\Gamma_{\chi_{c0}}$ of 21%.

Systematic errors on the branching fractions, which arise from the uncertainties in $\Gamma_{\chi_{c2}}$, the mass resolution, the choice of the background function, the efficiency determination, and the choices of the selection criteria are 11.5%, 12.6%, 12.1% and 14.7% for $B(\chi_{c0} \to \pi^+\pi^-)$, $B(\chi_{c0} \to K^+K^-)$, $B(\chi_{c2} \to \pi^+\pi^-)$ and $B(\chi_{c2} \to K^+K^-)$, respectively. There are overall errors caused by the uncertainty of the total number of $\psi(2S)$ events and the uncertainties in the $\psi(2S) \to \gamma\chi_{cJ}$ branching fractions. Adding these errors in quadrature gives total relative systematic errors of 15%, 16%, 16% and 18%, respectively, for $B(\chi_{c0} \to \pi^+\pi^-)$, $B(\chi_{c0} \to K^+K^-)$, $B(\chi_{c2} \to \pi^+\pi^-)$ and $B(\chi_{c2} \to K^+K^-)$.

For $B(\chi_{cJ} \to p\bar{p})$, sources of systematic errors include those listed above plus that associated with the assumption of an isotropic angular distribution for $\chi_{cJ} \to p\bar{p}$ decays. Adding all of the errors in quadrature gives relative systematic errors of 33%, 67% and 56% for the χ_{c0}, χ_{c1} and χ_{c2} states, respectively.

In summary, we obtain the total width of the χ_{c0} to be

$$\Gamma_{\chi_{c0}} = 14.3 \pm 2.0 \pm 3.0 \text{ MeV},$$

where the first error is statistical and the second is systematic. The results of the branching fractions of χ_{cJ} decays are listed in Table I and Table II. The measured width for the χ_{c0} is consistent with but substantially more precise than the previous measurement [2] (the uncertainty is reduced from 40% to 25%). The calculations involving new factorization schemes with high order QCD corrections [9,10] are in good agreement with our measurement.

Our branching fraction for $\chi_{c0} \to p\bar{p}$ is the first measurement for this decay, and is compatible with the previous upper bound [4]. Our results for $\chi_{c1,2} \to p\bar{p}$ decays, although statistically limited, are consistent, within errors, with the values determined from studies of charmonium states formed directly in $p\bar{p}$ annihilation [11]. The calculations from different models [12,13] give much smaller values of $\Gamma(\chi_{c0} \to p\bar{p})$ than our result. Our measurements on $\chi_{c0} \to p\bar{p}$, together with the results on $\eta_c \to p\bar{p}$ [7], indicate that the decay width of the helicity violating processes $\chi_{c0} \to p\bar{p}$ and $\eta_c \to p\bar{p}$ are much larger than those of $\chi_{c1,2} \to p\bar{p}$ decays which are allowed by the helicity conservation rule [3].

Finally, our branching fractions for $\chi_{c0,2}$ decays into $\pi^+\pi^-$ and K^+K^- are somewhat lower than the world average [7]. The recent calculations [14], which include color-octet contributions in exclusive χ_{cJ} decays, are in generally good agreement with our measurements. Using our results and canceling out the common errors in the branching fractions, we get the ratios of the branching fractions of $\frac{B(\chi_{c0}\to\pi^+\pi^-)}{B(\chi_{c0}\to K^+K^-)} = 0.82\pm0.15$ and $\frac{B(\chi_{c2}\to\pi^+\pi^-)}{B(\chi_{c2}\to K^+K^-)} = 1.88\pm0.51$.

We would like to thank the staffs of the BEPC accelerator and the Computing Center at the Institute of High Energy Physics, Beijing, for their outstanding scientific efforts. The work of the BES Collaboration was supported in part by the National Natural Science Foundation of China under Contract No. 19290400 and the Chinese Academy of Sciences under

contract No. KJ85 (IHEP), and by the Department of Energy under Contract Nos. DE-FG03-92ER40701 (Caltech), DE-FG03-93ER40788 (Colorado State University), DE-AC03-76SF00515 (SLAC), DE-FG03-91ER40679 (UC Irvine), DE-FG03-94ER40833 (U Hawaii), DE-FG03-95ER40925 (UT Dallas).

REFERENCES

† Deceased.

[1] G. T. Bodwin, E. Braaten and G. P. Lepage, Phys. Rev. **D51** (1995) 1125.

[2] J. E. Gaiser *et al.*, Phys. Rev. **D34** (1986) 711.

[3] S. J. Brodsky and G. P. Lepage, Phys. Rev. **D24** (1981) 2848.

[4] R. Brandelik *et al.*, Nucl. Phys. **B160** (1979) 426.

[5] V. L. Chernyak and A. R. Zhitnitsky, Phys. Rep. **112** (1984) 173.

[6] BES Collab., Nucl. Instr. Meth. **A344** (1994) 319.

[7] Particle Data Group, Phys. Rev. **D54** part I (1996) and references therein.

[8] P. K. Kabir and A. J. G. Hey, Phys. Rev. **D13** (1976) 3161.

[9] Han-Wen Huang and Kuang-Ta Chao, Phys. Rev. **D54** (1996) 6850.

[10] M. L. Mangano and A. Petrelli, Phys. Lett. **B352** (1995) 445; A. Petrelli, Phys. Lett. **B380** (1996) 159.

[11] T. A. Armstrong *et al.*, Nucl. Phys. **B373** (1992) 35; C. Baglin *et al.*, Phys. Lett. **B172** (1986) 455;

[12] M. Anselmino, R. Cancelliere and F. Murgia, Phys. Rev. **D46** (1992) 5049.

[13] M. Anselmino, F. Carnso and S. Forte, Phys. Rev. **D44** (1991) 1438.

[14] J. Bolz, P. Kroll and G. A. Schuler, Phys. Lett. **B392**, (1997) 198.

FIGURES



FIG. 1. The $\pi^+\pi^-$ mass distribution for selected $\psi(2S) \to \gamma\pi^+\pi^-$ events.



FIG. 2. The K^+K^- mass distribution for selected $\psi(2S) \to \gamma K^+K^-$ events.



FIG. 3. The $p\bar{p}$ mass distribution for selected $\psi(2S) \to \gamma p\bar{p}$ events.

TABLES

TABLE I. Branching fractions of $\chi_{cJ} \to 0^-0^-$

decay mode	N^{obs}	BR($\times 10^{-3}$)	PDG [7] (10^{-3})
$\chi_{c0} \to \pi^+\pi^-$	720 ± 32	$4.79 \pm 0.27 \pm 0.73$	7.5 ± 2.1
$\chi_{c0} \to K^+K^-$	774 ± 38	$5.82 \pm 0.36 \pm 0.94$	7.1 ± 2.4
$\chi_{c2} \to \pi^+\pi^-$	185 ± 16	$1.52 \pm 0.15 \pm 0.24$	1.9 ± 1.0
$\chi_{c2} \to K^+K^-$	115 ± 13	$0.81 \pm 0.14 \pm 0.14$	1.5 ± 1.1

TABLE II. Branching ratios of $\chi_{cJ} \to p\bar{p}$

decay mode	N^{obs}	BR($\times 10^{-5}$)	PDG [7] (10^{-5})
$\chi_{c0} \to p\bar{p}$	15.2 ± 4.1	$16.3 \pm 4.4 \pm 5.4$	< 90
$\chi_{c1} \to p\bar{p}$	4.2 ± 2.2	$4.3 \pm 2.3 \pm 2.9$	8.6 ± 1.2
$\chi_{c2} \to p\bar{p}$	4.7 ± 2.5	$5.9 \pm 3.1 \pm 3.3$	10.0 ± 1.0